

Terahertz Pioneers

A Series of Interviews With Significant Contributors to Terahertz Science and Technology

AS A TRIBUTE to individuals who have contributed significantly, and over many years, to the terahertz community, and as a guide and inspiration for those who are just beginning their professional association with this field of study, these transactions have included, on a regular basis, a series of biographical interviews with technical researchers who have appreciably impacted the THz community in a positive manner. In order to go beyond a strict technical review and to take better advantage of the information and commentary only available through a direct discussion, these articles take on a less formal style than the research articles that can be found within the remaining pages of the transactions. The Editor-in-Chief has taken some leeway in this regard, for the benefit of communicating more fully the character, experiences, and historic circumstances that have shaped our community and set the directions for our collective research. As a further means of assuring that the true flavor and circumstances of the contributions are expressed in the text, all of the articles are compiled after a face-to-face interview. The final text is shared with, and often helped considerably, by comments from the subject of the article.

This month we travel to Japan to speak with Dr. Hiromasa Ito,¹ a longterm affiliate of the Research Institute of Electrical Communication (RIEC), Tohoku University, Sendai, and a former Team Leader in Terahertz Photonics at Japan's premier science research institution, RIKEN (Institute of Physical and Chemical Research). Professor Ito is a link in a long chain of noted THz advocates and researchers at Tohoku.

This chain began with Jun-Ichi Nishizawa² who according to Ito, "predicted the possibility of using the resonance of phonons and molecular vibrations to generate THz electromagnetic radiation"³ in 1963. Nishizawa came to RIEC in 1953, and established

the RIKEN Center for Advanced Photonics in Sendai, Japan, near the site of the University, in 1990. Nishizawa was also the team leader for the first major THz program in Japan (Nishizawa Terahertz Purojekuto), which he started in 1987 at Tohoku.

The THz tradition at Tohoku continued with Professor Koji Mizuno⁴, who arrived in Sendai in 1968, and became RIKEN's first THz focused Team Leader in 1990. The reigns were handed off to Hiromasa Ito in 1998, and the tradition continues under Dr. Haroaki Minamide, who took over from Ito in 2010.

In 2005, RIKEN Sendai established an entire THz research group, composed of three THz teams—Tera-Photonics, Terahertz Sensing, and Terahertz Quantum Devices. Dr. Katsumi Midorikawa, who has since been promoted to Director of the Center for Advanced Photonics, RIKEN, originally led the group. Professor Chiko Otani has now taken over as Group Leader.

Professor Ito is best known for his development of the periodically poled lithium niobate and DAST crystal-based THz sources that have done much to open up the THz gap. As you will read, however, he is also responsible for a variety of laser and nonlinear optics devices that have found application in other frequency regimes. A competent theorist, and a gifted experimentalist, Ito was forced to come up with most of his device fabrication and test equipment by building it himself. He apparently has not suffered from the experience, as there are many in the world who would like to have the products of his research.

Ito is also a strong believer in university/industry teaming and has himself volunteered to serve this cause as leader of a technology transfer initiative, director of a technology incubator program and most recently, program officer for a 10 year collaborative university/industry THz technology development effort sponsored by the Japan Science and Technology agency.

Professor Ito has spent more than 50 years as a productive and innovative researcher and has had a great influence on the direction and success of terahertz development in Japan, as well as worldwide. I hope you will enjoy his story and take away something useful from his scientific pathway to success.

PETER H. SIEGEL, *Editor-in-Chief*
California Institute of Technology
Pasadena, CA 91125 USA

Digital Object Identifier 10.1109/TTHZ.2014.2376055

¹I sat down with Professor Hiromasa Ito in an office of a scenically situated building—which serves as the Sendai (Japan) site for RIKEN (Institute of Physical and Chemical Research)—atop a hill overlooking the countryside, on a wonderfully sunny, but crisp, day this past Fall. This humble but steadfast supporter of THz development is part of a long line of notables at the Tohoku University and its RIEC (Research Institute of Electrical Communication) going back to the very beginnings of interest in this frequency regime in the early 1970's. When his story is added to others in this series of articles, it completes a picture that places Japan in the forefront of THz development, both then and now.

²Nishizawa came to RIEC at Tohoku in 1953, and stayed on the faculty until 1990. Afterwards, he became the President of Tohoku and later, of Iwate Prefectural University and Tokyo Metropolitan University. The remarks attributed to Nishizawa come from a paper he wrote (and is often cited) that appeared in *Denshi-Kagaku (Electron. Sci.)*, vol. 14, p. 17, 1963. An interview and THz Pioneer article on Nishizawa is planned for these Transactions in spring 2015.

³Page 2173 from: Kazuhiro Imai, Kodo Kawase, Minamide Hiroaki, and Hiromasa Ito, "A chromatically injection-seeded terahertz-wave parametric generator," *Opt. Lett.*, vol. 27, no. 24, pp. 2173–2175, 2002.

⁴Koji Mizuno is also the subject of an article within this series: P.H. Siegel, "THz Pioneer Koji Mizuno: 50 Years in Submillimeter-Waves: From Otaku to Sensei," *IEEE Trans. THz Sci. Technol.*, vol. 3, no. 2, pp. 129–133, Mar. 2013.

Terahertz Pioneer: Hiromasa Ito

“Generating THz Energy is Crystal Clear”

Peter H. Siegel, *Fellow, IEEE*

HIROMASA ITO¹ grew up wanting to be a scientist like his father before him. The fifth of six children, he was raised in the aftermath of World War II, and like others who have been the subject of these articles, the impact of the war on Ito's life was more than an historic footnote. Ito's father, Yoji Ito² was a prominent member of the Japanese Navy (a Commander) and was extremely active in the development of Japan's first radar instrumentation in the early 1940's. Educated at the Technische Hochschule Dresden, Germany, under Heinrich Barkhausen,³ Yoji worked on high power magnetrons, including the infamous Ku-Go (a microwave variation of Tesla's invisible death ray concept [1]). After the war he was severely restricted in his employment, but he managed to start a company, Koden Electronics, for the peacetime use of his Naval radio technology⁴. Koden Electronics, Tokyo, Japan, still exists today and is known for marine navigation and fish finders.

At the age of 10, Hiromasa's mentoring was left in the hands of a close friend and scientific colleague of Yoji's, Yasushi Watanabe.⁵ Hiromasa's older siblings all left their small agricultural township outside Tokyo as soon as they were of age. His oldest brother went to Tohoku University to serve in Watanabe's research group. Hiromasa left home to attend an excellent Junior and Senior high school at University of Tsukuba, Otsuka, Tokyo, which specialized in science. One of the teachers there encouraged him to study English, and after graduating, he followed his oldest brother, and enrolled at Tohoku University in Sendai in 1962.

Manuscript received November 24, 2014; accepted November 25, 2014. Date of publication December 19, 2014; date of current version January 14, 2015.

The author is with Department of Electrical Engineering, California Institute of Technology, Pasadena, CA 91125 USA (e-mail: phs@caltech.edu).

Digital Object Identifier 10.1109/THZ.2014.2376054

¹I sat down with Professor Hiromasa Ito in an office of a scenically situated building—which serves as the Sendai (Japan) site for RIKEN (Institute of Physical and Chemical Research)—atop a hill overlooking the countryside, on a wonderfully sunny, but crisp, day this past Fall. This humble but steadfast supporter of THz development is part of a long line of notables at the Tohoku University and its RIEC (Research Institute of Electrical Communication) going back to the very beginnings of interest in this frequency regime in the early 1970's. When his story is added to others in this series of articles, it completes a picture that places Japan in the forefront of THz development, both then and now.

²For a short biography of Yoji Ito, see: http://en.wikipedia.org/wiki/Yoji_Ito

³Barkhausen was the inventor of the Barkhausen-Kurz microwave oscillator, a precursor of the klystron.

⁴Yoji developed a critically useful marine radio direction sensor as an aid to navigation. <https://www.koden-electronics.co.jp/eng/about/corp-e-story.html>

⁵Watanabe was a noted electrical engineer, department chair at Tohoku University and later, President of Shizuoka University. He remained a close family friend of the Itos, serving as Ceremonial Go-Between (a very honored family role) in the marriages of all 4 sons of Yoji, including Hiromasa. <http://www.shmj.or.jp/english/pioneers/pnr23.html>.



HIROMASA ITO

Hiromasa was interested in atomic physics, but Watanabe recommended electronics. Ito was to make an interesting compromise. In his third year at Tohoku, he found life-long mentor and colleague, Humio Inaba, then a new faculty member at the Research Institute for Electrical Communication (RIEC) who had recently come back from a research visit at Stanford University, CA, USA, with knowledge of lasers and masers. Inaba was a physicist, but had a strong background in electronics, and was working in the Tohoku antennas group led by Hidenari Uchida. Uchida believed that radio antennas and masers would play a key role in space communications. It was due to Inaba's contributions to the Ibaraki Earth station in Takahagi, Japan that it was ready to receive the first transmission test of a transpacific satellite broadcast between the USA and Japan via Relay 1 on November 23, 1963. The transmission was planned to be an address by U.S. President John F. Kennedy, but instead was an historic broadcast containing news of Kennedy's assassination in Dallas, TX, USA, on Nov. 22, 1963.

Ito began working with Inaba on Ruby lasers and the earliest forms of laser radar (LIDAR) [2]. His Master's degree focused on mode locking [3] and other characteristics of newly invented CO₂ lasers. This included the first observation of beam spreading in liquids [4], [5]. Everything had to be fabricated and assembled locally. Ito became proficient at power electronics, glass blowing, optical polishing and coatings, crystal growth and thin film deposition.

After completing his dissertation in 1971 and joining RIEC as a research associate, he began to become interested in optical conversion, and especially in optical parametric oscillators (OPOs). Working with Inaba, he obtained parametric fluorescence generation from 4580 to 6100 Å in lithium formate monohydrate pumped by a UV nitrogen gas laser [6]. This was followed by experimental and theoretical work on sodium formate and lithium–sodium formate crystals with high second harmonic generation efficiency (for reaching into the UV) as well as broad OPO tunability (from the optical out to far IR wavelengths) when pumped in the UV with Nd:YAG lasers [7]–[9].

In order to expand upon his nonlinear optical experiments, Ito needed to acquire the pump lasers, which were not available in Japan at the time. When Nd:YAG lasers first became commercially available in the early 1970s, his lab hosted a visit from noted Stanford Professor and laser pioneer, Robert Byer (2013 American Physical Society President), who was helping promote Nd:YAG lasers for a U.S. company. Ito became so involved, that he helped a Japanese import/export company add the laser as a product for research laboratories around the country. He also linked up with Byer, who would later (1975/1976) host him (Ito) at Stanford University, and who remained a lifelong friend and colleague.

In 1974, Ito had the first verification of phase-matched second-harmonic generation in a thin film nonlinear waveguide structure composed of ZnS deposited on lithium niobate [10]–[12]. This work led to the development of a much more efficient 4 layer structure composed of linear and nonlinear polarized materials which he managed to fabricate himself at RIEC [13], [14]. Continuing with his interests in optical waveguides and strong nonlinear crystal interactions, Ito produced an optical switch [15] and directional couplers [16] based on LiNbO₃, as well as an early concept for an infrared difference frequency generator based on semiconductor diode laser mixing in his nonlinear optical waveguide structures [17]. He also observed picosecond [18] and later sub-picosecond [19]–[21] pulsing with very high repetition rate in AlGaAs laser diodes, later to become an important component concept for optical communications.

In 1982, Ito moved from staff researcher to Associate Professor at Tohoku University and started working on a series of novel semiconductor laser devices. He was only able to fabricate the devices by locally assembling III-V semiconductor material MOCVD (metal–organic chemical-vapor deposition), LPE (liquid phase epitaxy), and RIE (reactive ion etch) systems [22], [23]. Ito used the III-V semiconductor fabrication equipment to invent a new type of vertical cavity surface emitting light structure that he termed the coaxial transverse junction (CTJ) laser/LED [24], [25]. He helped transition this technology, at an early stage, to the RICOH Co. laboratory in Sendai for the development of their optical heads used in print/copy machines. RICOH turned the invention into a commercial process, and is now using this technology quite successfully in many of their products.

Ito's sojourn into the THz domain had its origins with his work on periodically poled (domains of alternating polarization) grating structures fabricated on LiNiO₃ in 1988 [26]. Quasi-phase matching conditions for amplification and efficient signal beam output within the optical waveguide were realized by a poled grating deposited through a simple room temperature scanning electron beam technique [27]. Efficient

second harmonic output was also obtained through a Cerenkov quasi-phase matching approach controlled by a periodic grating, where the output beam appears at a Cerenkov angle established by the grating properties [28]. These developments set the stage for a fortuitous occurrence that was to dramatically alter the direction of Ito's research.

In February 1990, a symposium was held at Tohoku by Jun-ichi Nishizawa⁶, then director of RIEC, and in charge of a specially funded Japanese Science and Technology agency program on Terahertz. The theme of the symposium was “bridging the gap between light and electromagnetic waves” [29]. Nishizawa was moderating a panel discussion when he posed a question to the audience as to what techniques could best join these two dramatically disparate photonic and electronic regimes. Scanning the crowd he pointed dramatically at Hiromasa Ito, who was quietly listening from a front row seat, and asked “What do you think about this?” It was a moment of *truth or dare!* Ito responded rather boldly, “I believe I can use nonlinear optics to bridge the gap.” *A new recruit had just entered the ranks of THz researchers in Japan!*

Two years later a young, and very determined graduate student knocked on the door. Kodo Kawase had just come from a meeting with Humio Inaba, where he had told Inaba that he wanted to work on the interaction of electromagnetic waves with biological systems for his dissertation. He was particularly excited about THz waves, because of a recent resurgence of interest in potential biological interactions originally proposed by Fröhlich [30], [31]. Inaba sent Kawase over to Ito, who when he (Ito) heard about the biological interest, and realized how difficult a topic this would be for a thesis project at RIEC, wisely suggested to Kawase that before he could explore any long wave electromagnetic interactions with tissue, he would need to have a strong and broadly tunable THz source. This argument convinced Kawase that working on sources with Ito would make a good start to his long term research goals.

THz generation through laser pumped crystals with strong nonlinear susceptibility [32], specifically lithium niobate [33], [34] had been demonstrated in the early as 1970's (see also [35]). However little improvements had been made on these systems since that time, and neither CW nor pulsed versions had been realized with sufficient efficiency and THz output power for the experiments Kawase was eager to perform. The recently demonstrated periodically poled crystalline structures Ito had been developing [36] just might prove to be the breakthrough that was required to turn these earlier demonstrations into solidly functioning instruments. So in 1992, Kawase set to work on the optically pumped THz source problem under Ito's supervision.

The first results from Kawase's and Ito's efforts to generate THz signals from their periodically poled lithium niobate structures involved difference frequency mixing in a variety of crystalline structures and appeared in 1994 [37]. Although the design was presented, there was too much THz absorption in the

⁶Nishizawa is an extremely prominent figure, and himself a pioneer in THz science. He was the leader of the Japanese Science and Technology agency's Exploratory Research for Advanced Technology Office (ERATAO) program on Terahertz between 1987–1992 and director of RIEC from 1988 to 1990. After this he became President of Tohoku University (1990–1996), President of Iwate Prefectural University (1998–2005) and President of Tokyo Metropolitan University (2005–2009). The IEEE issued an award in his honor, the Jun-ichi Nishizawa medal, in 2002. With a bit of luck we will have a Pioneer article on this notable figure in an upcoming issue of these transactions.

crystals, and the index of refraction of LiNbO_3 was too high at these wavelengths (> 5), to get more than a fluorescence signature. Kawase and Ito had an idea for reducing the THz absorption and output coupling mismatch by introducing a grating on top of the lithium niobate to steer the beam out of the crystal. The fabrication was tedious as the grating was introduced along the top propagation plane of the LiNbO_3 by using a diamond wafer saw to precisely groove the crystalline surface. The Q-switched Nd:YAG laser pump beam was directed at the ends of the crystal and the near infrared idler (allowed to exit the ends of the crystal through an appropriate antireflection coating) was resonated by a high Q external cavity. By design, the non-collinear phase matching conditions would shift the idler-to-pump propagation angle and the angle of the generated THz signal beam sufficiently such that all three wavelengths could be spatially separated. The grating further directed the THz beam so it exited the top of the crystal at a relatively steep angle compared to the pump and idler paths. For detecting the THz energy, Kawase and Ito set up both a helium cooled IR bolometer and a Schottky barrier diode detector supplied by colleague, Koji Mizuno [38], who also taught the duo useful techniques for performing sub-millimeter-wave measurements.

Just as the first experiments were getting underway in late 1995, Ito was recruited by the Japanese government to go on an extended trip to the U.S. and Europe as part of a program to stimulate technology transfer between government laboratories and industry partners. Kawase sent daily faxes to the hotels where Ito was staying after each day's measurements. Finally, in November of 1995 they achieved THz first light. By slightly changing the angle of the pump beam to the crystal axis (< 1 degree), the THz output could be tuned from 180 to 270 microns (1.1–1.66 THz). The output power from the grating coupler was 250 times greater than the energy that would have exited a Brewster angled edge cut in the crystal z-axis, and was in the mW range—far higher than any previous experiments. The published results appeared in April 1996 [39].

Over the next two years, Kawase, Ito, and RIEC colleagues realized higher power, broader tuning, tighter beam forming, and more compact packaging for the THz OPO [40]–[43]. The silicon prism coupler [41], an idea of Kawase's, allowed the THz beam to remain at a fixed angle relative to the crystal during frequency tuning, and enabled a fully packaged source to be assembled and eventually commercialized⁷.

In 1998, Ito was appointed Team Leader for Tera Photonics at the RIKEN (Institute of Physical and Chemical Research) Photodynamics Research Center in Sendai, where he replaced Koji Mizuno. By this stage, the groundbreaking experiments on high power THz generation from LiNbO_3 had led to an explosion of further development, with worldwide interest in this “reborn” and greatly improved technology.

Between 1998 and 2005, the period of Ito's first appointment as a RIKEN team leader, he, in conjunction with his colleagues and staff at Tohoku, published more than 70 journal articles, and hundreds of conference papers on terahertz sources and applications, only a selected group of which have been reproduced in the references [44]–[70]. Significant milestones include: injection seeding via a yttrium fiber laser to achieve narrow line width (100 MHz) [47], [48], [52], surface-emission difference frequency mixing in a periodically poled structure

[46], [55], grating tuned injection seeding to allow continuous rapidly tuned THz output without moving lenses or mirrors [56], continuous wave THz output [67], and THz generation from a two-dimensional PPLN (periodically poled lithium niobate) crystal [68]. These sources have opened up the THz bands for a wide range of applications from imaging, to spectroscopy, to the study of ultra-high field and nonlinear optical effects, and much, much more.

While improving upon lithium niobate structures, Ito was also working on new crystals with even higher nonlinearities for use as THz sources. An organic salt known as 4-N,N-dimethylamino-4'-N'-methyl-stilbazolium tosylate (DAST), invented by Tohoku's Hachiro Nakanishi [71], enabled Ito and colleagues to significantly improve the THz OPO efficiency and bandwidth [72]–[74]. DAST is still being grown at RIKEN, (I had a chance to see it up close while visiting), and although it has not yet been possible to achieve large, defect free crystals, DAST has been utilized in a wide variety of THz source and detection systems at Tohoku [75]–[81]. Since being popularized by Ito's team, especially Kawase [74], DAST is now seeing applications worldwide ([82], [83], for example).

As the PPLN OPO THz sources became more and more “user friendly” and portable, it was natural for Ito and coworkers to get involved in THz applications. Although most of these were pioneered and publicized by RIKEN Tera Photonics staff members, there were several standouts with Ito's involvement, including chemical signaturing [61], [62], near field imaging [65], [69] and imaging for a variety of commercial, medical, and security applications [66].

During all of the THz source development activities, Ito also managed to come up with an all solid-state acoustically modulated laser structure that could produce very precise continuously chirped combs—the frequency-shifted feedback laser (FSFL) [84]–[87]. This turned out to have many commercial applications including ranging [88], dispersion measurements [89], and communications [90]. It is most recently being used to perform precise 3D surface contouring at distances from 1–5 meters with accuracies in the micron range [91].

Ito retired from Tohoku University in 2007 after serving as director of RIEC for three years, but took up a second term as Team Leader at RIKEN from 2005 through to 2010. If you were under the impression that this “retirement” slowed him down, you would be dead wrong. Between 2005 and 2010 Ito plunged into THz devices and applications with a passion, publishing 40+ journal articles in the field, in addition to continued development of the frequency shifted feedback lasers, DAST and other nonlinear crystals, new low loss THz waveguide structures and new THz detectors based on pumped nonlinear crystal up-conversion processes [92]–[98]. He worked most closely with Hiroaki Minamide during this period, who became the Team Leader for the RIKEN group in 2010, and is carrying on both the work of Professor Ito, as well as many new and exciting projects involving THz generators and detectors based on nonlinear optical phenomena and techniques [99], [100].

After 2010, Professor Ito shifted over to Senior Scientist at RIKEN and in his “spare time” he took on roles as Director of the Japan Science and Technology (JST) agency *Innovation Plaza Miyagi*, where he was able to work towards promoting university and industry collaborations. He then transitioned into a program officer position at JST to shepherd collaborative university research into industrial demand. Ito continues to do re-

⁷The LiNbO_3 source concept was incorporated into the “Goemon-R” (Phluxi, Inc., Sendai, Japan).

search, to mentor students and colleagues, and to publish copiously ([101]–[108] to list just a few), as he celebrates his 53rd year at Tohoku. The last [109], just recently published in 2014, contains a nice concise summary of the work he and his colleagues performed at RIKEN.

It must be very satisfying indeed to think back to that day in February 1990, when his very public response to Sensei Jun-ichi Nishizawa about linking optics to electromagnetics was, “*I believe I can use nonlinear optics to bridge the gap.*” Clearly, Sensei Hiromasa Ito has now fulfilled that pledge.

ACKNOWLEDGMENT

The author would like to warmly thank Profs. K. Kawase and K. Mizuno for their help in setting up this interview and their kind reception in Sendai. Special thanks also go to Dr. H. Minamide, the current Team Leader of the Tera photonics group at RIKEN, for the use of his office, a wonderful tour of the RIKEN facilities, and a warm reception there.

REFERENCES

- [1] W. E. Grunden, “Secret weapons and World War II: Japan in the shadow of Big Science,” in *Electric Weapons: Radar and the Death Ray*. Manhattan, KS: Univ. Kansas Press, 2005, ch. 3, pp. 83–123.
- [2] H. Inaba, T. Kobayashi, T. Ichimura, M. Morishita, and H. Ito, “Fundamental study of operational performance of a laser radar system employing A-scope representation,” *Electron. Commun. Jpn.*, vol. 51-B, no. 9, pp. 36–44, 1968.
- [3] H. Ito and H. Inaba, “Self mode-locking of the transverse modes in the CO₂ laser oscillation,” *Opt. Commun.*, vol. 1, no. 2, pp. 61–63, 1969.
- [4] H. Inaba and H. Ito, “Observation of power-dependent distortion of an infrared beam at 10.6 μm from a CO₂ laser during propagation in liquids,” *IEEE J. Quantum Electron.*, vol. QE-4, no. 2, pp. 45–48, Feb. 1968.
- [5] H. Inaba and H. Ito, “Self-defocusing effect of infrared CO₂ laser beams induced thermally in liquids,” *IEEE J. Quantum Electron.*, vol. QE-4, no. 5, p. 351, May 1968.
- [6] H. Ito and H. Inaba, “Optical properties and UV N₂ laser-pumped parametric fluorescence in LiCOOH:H₂O,” *IEEE J. Quantum Electron.*, vol. QE-8, no. 6, p. 612, Jun. 1972.
- [7] H. Ito, H. Naito, and H. Inaba, “New phase-matchable nonlinear optical crystals of Formate family,” *IEEE J. Quantum Electron.*, vol. QE-9, no. 6, pp. 701–702, Jun. 1973.
- [8] H. Ito, H. Naito, and H. Inaba, “New phase-matchable nonlinear optical crystals of the Formate family,” *IEEE J. Quantum Electron.*, vol. QE-10, no. 2, pp. 247–252, Feb. 1974.
- [9] H. Ito, H. Naito, and H. Inaba, “Generalized study on angular dependence of induced second-order nonlinear optical polarizations and phase matching in biaxial crystals,” *J. Appl. Phys.*, vol. 46, no. 9, pp. 3992–3998, 1975.
- [10] H. Ito, N. Uesugi, and H. Inaba, “Phase-matched guided optical second harmonic wave generation in oriented ZnS polycrystalline thin-film waveguides,” *Appl. Phys. Lett.*, vol. 25, no. 7, pp. 385–387, 1974.
- [11] H. Ito and H. Inaba, “Phase-matched SHG in thin-film nonlinear optical-waveguide incorporated with substrate nonlinearity,” *IEEE J. Quantum Electron.*, vol. 11, no. 9, p. 862, Sep. 1975.
- [12] H. Ito and H. Inaba, “Phase-matched guided, optical second harmonic generation in nonlinear ZnS thin-film waveguide deposited on nonlinear LiNbO₃ substrate,” *Opt. Commun.*, vol. 15, no. 1, pp. 104–107, 1975.
- [13] H. Ito and H. Inaba, “Study on phase matching characteristics of optical second harmonic generation in nonlinear thin-film waveguides using a tunable parametric oscillator,” in *Tunable Lasers and Applications*. Berlin, Germany: Springer Verlag, 1976, pp. 353–360.
- [14] H. Ito and H. Inaba, “Efficient phase-matched second-harmonic generation method in four-layered optical-waveguide structure,” *Opt. Lett.*, vol. 2, no. 6, pp. 139–141, 1978.
- [15] H. Ito, Y. Ogawa, and H. Inaba, “Integrated bistable optical device using Mach-Zehnder interferometric optical waveguide,” *Electron. Lett.*, vol. 15, no. 10, pp. 283–285, 1979.
- [16] H. Ito, Y. Ogawa, K. Makita, and H. Inaba, “Integrated bistable optical multivibrator using electro-optically controlled directional coupler switches,” *Electron. Lett.*, vol. 15, no. 24, pp. 791–793, 1979.
- [17] H. Ito and H. Inaba, “Integrated coherent infrared generator operated at room temperature using nonlinear optical waveguide structure,” *Proc. IEEE*, vol. 67, no. 10, pp. 1455–1456, Oct. 1979.
- [18] H. Ito, H. Yokoyama, S. Murata, and H. Inaba, “Picosecond optical pulse generation from an R.F. modulated AlGaAs D.H. diode laser,” *Electron. Lett.*, vol. 15, no. 23, pp. 738–740, 1979.
- [19] H. Yokoyama, H. Ito, and H. Inaba, “Generation of subpicosecond coherent optical pulses by passive-mode locking of an AlGaAs diode-laser,” *Appl. Phys. Lett.*, vol. 40, no. 2, pp. 105–107, 1982.
- [20] Y. Tada, H. Yokoyama, H. Ito, and H. Inaba, “Ultrashort optical pulse generation from microwave modulated AlGaAs diode-laser with self-coupled rod resonator,” *Opt. Commun.*, vol. 47, no. 3, pp. 187–189, 1983.
- [21] H. Ito, N. Onodera, K. Gen-ei, and H. Inaba, “Self-Q-switched picosecond optical pulse generation with tandem-type AlGaAs TJS laser,” *Electron. Lett.*, vol. 17, no. 1, pp. 15–17, 1981.
- [22] H. Yamada, H. Ito, and H. Inaba, “Microprocessing of GaAs cylindrical columns for integrated optical-device fabrication by Cl₂-Ar reactive ion etching,” *Electron. Lett.*, vol. 20, no. 14, pp. 591–592, 1984.
- [23] H. Yamada, H. Ito, and H. Inaba, “Anisotropic reactive ion etching technique of GaAs and AlGaAs materials for integrated optical-device fabrication,” *J. Vac. Sci. Technol. B*, vol. 3, no. 3, pp. 884–888, 1985.
- [24] H. Ito, N. Komagata, H. Yamada, and H. Inaba, “Novel structure of laser diode and light-emitting diode realized by coaxial transverse junction (CTJ),” *Electron. Lett.*, vol. 20, no. 14, pp. 577–579, 1984.
- [25] H. Yamada, H. Watanabe, H. Ito, and H. Inaba, “Surface-emitting GaAs light-emitting diode-laser diode with modified coaxial transverse junction (CTJ) structure,” *Electron. Lett.*, vol. 24, no. 2, pp. 77–78, Jan. 1988.
- [26] Y. H. Zhang, H. Ito, and H. Inaba, “Experiments on nonlinear optical waveguide with periodic ferroelectric domain inversion,” in *Annu. Meeting Jpn. Soc. Appl. Phys. (JSAP)*, 1988, 7a-ZD-9.
- [27] H. Ito, C. Takyu, and H. Inaba, “Fabrication of periodic domain grating in LiNbO₃ by electron-beam writing for application of nonlinear optical processes,” *Electron. Lett.*, vol. 27, no. 14, pp. 1221–1222, Jul. 1991.
- [28] H. Ito, T. Fujiwara, and C. Takyu, “Cerenkov-type second harmonic generation from LiNbO₃ waveguide with periodic domain grating,” *Opt. Commun.*, vol. 99, pp. 237–240, 1993.
- [29] J. Nishizawa, “Boundaries between light and electromagnetic waves,” in *RIEC Symp.*, Sendai, Japan, Feb. 1–2, 1990.
- [30] H. Fröhlich, *Biological Coherence and Response to External Stimuli*. Berlin, Germany: Springer-Verlag, 1988.
- [31] P. H. Siegel, “THz pioneer Fritz Keilmann: ‘RF biophysics: From strong field to near field’,” *IEEE Trans. THz Sci. Technol.*, vol. 3, no. 5, pp. 505–513, Sep. 2013.
- [32] J. R. Morris and Y. R. Shen, “Far-Infrared generation by picosecond pulses in electro-optical materials,” *Opt. Commun.*, vol. 3, no. 2, pp. 81–84, 1971.
- [33] K. H. Yang, P. L. Richards, and Y. R. Shen, “Generation of far-infrared radiation by picosecond light pulses in LiNbO₃,” *Appl. Phys. Lett.*, vol. 19, pp. 320–322, Nov. 1971.
- [34] M. A. Piestrup, R. N. Fleming, and R. H. Pantell, “Continuously tunable submillimeter wave source,” *Appl. Phys. Lett.*, vol. 26, no. 8, pp. 418–420, 1975.
- [35] P. H. Siegel, “Terahertz pioneer: Michael Bass ‘The THz light at the end of the tunnel’,” *IEEE Trans. THz Sci. Technol.*, vol. 4, no. 4, pp. 409–417, Jul. 2014.
- [36] H. Ito, “Nonlinear optics using periodic domain inverted structures, Nonlinear optics,” in *Proc. Int. Conf. Optically Nonlinear Organic Mater. Appl.*, T. Kobayashi, Ed., May 17–20, 1994, pp. 327–331.
- [37] K. Kawase and H. Ito, “Submillimeter generation using periodic domain reversals,” in *Proc. Int. Conf. Optically Nonlinear Organic Mater. Appl.*, T. Kobayashi, Ed., May 17–20, 1994, pp. 225–229.
- [38] P. H. Siegel, “THz pioneer Koji Mizuno: ‘50 years in submillimeter-waves: From Otaku to Sensei’,” *IEEE Trans. THz Sci. Technol.*, vol. 3, no. 2, pp. 129–133, Mar. 2013.
- [39] M. Sato, T. Taniuchi, and H. Ito, “Coherent tunable THz-wave generation from LiNbO₃ with monolithic grating coupler,” *Appl. Phys. Lett.*, vol. 68, no. 18, pp. 2483–2485, Apr. 1996.
- [40] K. Kawase, M. Sato, T. Taniuchi, and H. Ito, “Characteristics of THz-wave radiation using a monolithic grating coupler on a LiNbO₃ crystal,” *Int. J. Infrared Millim. Waves*, vol. 17, no. 11, pp. 1839–1849, Nov. 1996.
- [41] K. Kawase, M. Sato, K. Nakamura, T. Taniuchi, and H. Ito, “Unidirectional radiation of widely tunable THz wave using a prism coupler under noncollinear phase matching condition,” *Appl. Phys. Lett.*, vol. 71, no. 6, pp. 753–755, Aug. 1997.

- [42] H. Ito, K. Kawase, and J. Shikata, "Widely tunable THz-wave generation by nonlinear optics," *IEICE Trans. Electron.*, vol. E81C, no. 2, pp. 264–268, Feb. 1998.
- [43] J.-I. Shikata, M. Sato, T. Taniuchi, and H. Ito, "Enhancement of terahertz-wave output from LiNbO₃ optical parametric oscillators by cryogenic cooling," *Opt. Lett.*, vol. 24, no. 4, pp. 202–204, 1999.
- [44] A. Morikawa, K. Kawase, J. Shikata, T. Taniuchi, and H. Ito, "Parametric THz-wave generation using trapezoidal LiNbO₃," in *Proc. SPIE THz Spectrosc. Appl. II*, J. M. Chamberlain, Ed., 1999, pp. 302–310.
- [45] J. Shikata, K. Kawase, K. Karino, T. Taniuchi, and H. Ito, "Tunable terahertz-wave parametric oscillators using LiNbO₃ and MgO:LiNbO₃ crystals," *IEEE Trans. Microw. Theory Tech.*, vol. 48, no. 4, pp. 653–661, Apr. 2000.
- [46] Y. H. Avetisyan, K. Kawase, and H. Ito, "Surface-emitted difference frequency generation in non-ferroelectric materials," in *Proc. SPIE THz and GHz Electron. Photon. II*, 2000, vol. 4111, pp. 382–390.
- [47] K. Imai, K. Kawase, J.-I. Shikata, H. Minamide, and H. Ito, "Injection-seeded terahertz-wave parametric oscillator," *Appl. Phys. Lett.*, vol. 78, no. 8, pp. 1026–1028, 2001.
- [48] K. Kawase, J.-I. Shikata, K. Imai, and H. Ito, "Transform-limited, narrow-linewidth, terahertz-wave parametric generator," *Appl. Phys. Lett.*, vol. 78, no. 19, pp. 2819–2821, 2001.
- [49] K. Kawase, J. Shikata, H. Minamide, K. Imai, and H. Ito, "Arrayed silicon prism coupler for a THz-wave parametric oscillator," *Applied Opt.*, vol. 40, no. 9, pp. 1423–1426, 2001.
- [50] K. Imai, K. Kawase, and H. Ito, "A frequency-agile terahertz-wave parametric oscillator," *Opt. Express*, vol. 8, no. 13, pp. 699–704, 2001.
- [51] A. Sato, K. Kawase, H. Minamide, S. Wada, and H. Ito, "Tabletop terahertz-wave parametric generator using a compact, diode-pumped Nd:YAG laser," *Rev. Sci. Instrum.*, vol. 72, no. 9, pp. 3501–3504, 2001.
- [52] K. Kawase, H. Minamide, K. Imai, J.-I. Shikata, and H. Ito, "Injection-seeded terahertz-wave parametric generator with wide tenability," *Appl. Phys. Lett.*, vol. 80, no. 2, pp. 195–197, 2002.
- [53] K. Kawase, J. Shikata, and H. Ito, "Terahertz-wave parametric source," *J. Phys. D—Appl. Phys. Topical Rev.*, vol. 35, no. 3, pp. R1–R14, Feb. 2002.
- [54] A. Sato, K. Imai, K. Kawase, H. Minamide, S. Wada, and H. Ito, "Narrow-linewidth operation of a compact THz-wave parametric generator system," *Opt. Commun.*, vol. 207, no. 1–6, pp. 353–359, 2002.
- [55] Y. Sasaki, A. Yuri, K. Kawase, and H. Ito, "Terahertz-wave surface-emitted difference frequency generation in slant-stripe-type periodically poled LiNbO₃ crystal," *Appl. Phys. Lett.*, vol. 81, no. 18, pp. 3323–3325, 2002.
- [56] K. Imai, K. Kawase, M. Hiroaki, and H. Ito, "Achromatically injection-seeded terahertz-wave parametric generator," *Opt. Lett.*, vol. 27, no. 24, pp. 2173–2175, 2002.
- [57] K. Imai, S. Sugawara, J. Shikata, K. Kawase, H. Minamide, and H. Ito, "The effect of injection seeding on terahertz parametric oscillation," *Electron. Commun. in Japan, Part II—Electron.*, vol. 86, no. 1, pp. 26–35, 2003.
- [58] K. Kawase, J. Shikata, and H. Ito, "Narrow-linewidth tunable terahertz-wave sources using nonlinear optics," *Solid-State Mid-Infrared Laser Sources*, vol. 89, pp. 397–423, 2003.
- [59] J. Shikata, K. Kawase, H. Ito, and R. Wako, "The generation and linewidth control of terahertz waves by parametric processes," *Electron. Commun. in Japan, Part II—Electron.*, vol. 86, no. 5, pp. 52–65, 2003.
- [60] T. Suhara, Y. Avetisyan, and H. Ito, "Theoretical analysis of laterally emitting terahertz-wave generation by difference-frequency generation in channel waveguides," *IEEE J. Quantum Electron.*, vol. 39, no. 1, pp. 166–171, Jan. 2003.
- [61] Y. Watanabe, K. Kawase, T. Ikari, H. Ito, Y. Ishikawa, and H. Minamide, "Component spatial pattern analysis of chemicals using terahertz spectroscopic imaging," *Appl. Phys. Lett.*, vol. 83, no. 4, pp. 800–802, Jul. 2003.
- [62] Y. Watanabe, K. Kawase, T. Ikari, H. Ito, Y. Ishikawa, and H. Minamide, "Component analysis of chemical mixtures using terahertz spectroscopic imaging," *Opt. Commun.*, vol. 234, pp. 125–129, 2004.
- [63] Y. Sasaki, H. Yokoyama, and H. Ito, "Dual-wavelength optical-pulse source based on diode lasers for high-repetition-rate, narrow-bandwidth terahertz-wave generation," *Opt. Express*, vol. 12, no. 14, pp. 3066–3071, 2004.
- [64] T. Hidaka, H. Minamide, H. Ito, J. Nishizawa, K. Tamura, and S. Ichikawa, "Ferroelectric PVDF cladding terahertz waveguide," *J. Lightw. Technol.*, vol. 23, no. 8, pp. 2469–2473, Aug. 2005.
- [65] K. Ishihara, T. Ikari, H. Minamide, J. Shikata, K. Ohashi, H. Yokoyama, and H. Ito, "Terahertz near-field imaging using enhanced transmission through a single subwavelength aperture," *Jpn. J. Appl. Phys.*, vol. 44, no. 28–32, pt. 2, pp. L929–L931, 2005.
- [66] K. Kawase, Y. Ogawa, H. Minamide, and H. Ito, "Terahertz parametric sources and imaging applications," *Semicond. Sci. Technol.*, vol. 20, no. 7, pp. S258–S265, Jul. 2005.
- [67] Y. Sasaki, H. Yokoyama, and H. Ito, "Surface-emitted continuous-wave terahertz radiation using periodically poled lithium niobate," *Electron. Lett.*, vol. 41, no. 12, pp. 712–713, Jun. 2005.
- [68] Y. Sasaki, Y. Avetisyan, H. Yokoyama, and H. Ito, "Surface-emitted terahertz-wave difference frequency generation in two-dimensional periodically poled lithium niobate," *Opt. Lett.*, vol. 30, no. 21, pp. 2927–2929, 2005.
- [69] K. Ishihara, K. Ohashi, T. Ikari, H. Minamide, H. Yokoyama, J. Shikata, and H. Ito, "Terahertz-wave near-field imaging with sub-wavelength resolution using surface-wave-assisted bow-tie aperture," *Appl. Phys. Lett.*, vol. 89, no. 20, p. 201120, Nov. 2006.
- [70] R. X. Guo, K. Akiyama, H. Minamide, and H. Ito, "All-solid-state, narrow linewidth, wavelength-agile terahertz-wave generator," *Appl. Phys. Lett.*, vol. 88, no. 9, p. 091120, Feb. 2006.
- [71] S. Okada, H. Matsuda, H. Nakanishi, M. Kato, and R. Muramatsu, "Organic material for non-linear optical use," Japan Patent Appl. 61-192404, Feb. 29, 1988, Publication number 63-048265.
- [72] K. Kawase, M. Mizuno, S. Sohma, H. Takahashi, T. Taniuchi, Y. Urata, S. Wada, H. Tashiro, and H. Ito, "Difference-frequency terahertz-wave generation from 4-dimethylamino-N-methyl-4-stilbazolium-tosylate by use of an electronically tuned Ti:sapphire laser," *Opt. Lett.*, vol. 24, no. 15, pp. 1065–1067, 1999.
- [73] S. Sohma, H. Takahashi, T. Taniuchi, and H. Ito, "Organic nonlinear optical crystal DAST growth and its device applications," *Chem. Phys.*, vol. 245, no. 1–3, pp. 359–364, 1999.
- [74] K. Kawase, T. Hatanaka, H. Takahashi, K. Nakamura, T. Taniuchi, and H. Ito, "Tunable THz-wave generation from DAST crystal using dual signal-wave parametric oscillation of periodically poled lithium niobate," *Opt. Lett.*, vol. 25, pp. 1714–1716, 2000.
- [75] K. Suizu, K. Miyamoto, T. Yamashita, and H. Ito, "High-power terahertz-wave generation using DAST crystal and detection using mid-infrared power meter," *Opt. Lett.*, vol. 32, no. 19, pp. 2885–2887, Oct. 2007.
- [76] T. Taniuchi, J. Shikata, and H. Ito, "Tunable terahertz-wave generation in DAST crystal with dual-wavelength KTP optical parametric oscillator," *Electron. Lett.*, vol. 36, no. 16, pp. 1414–1416, Aug. 2000.
- [77] Y. Namba, S. Koyama, H. Ito, Y. M. Son, and N. Ohsaki, "Optical fabrication of organic nonlinear DAST crystals for high frequency modulation," in *Proc. SPIE 4829, 19th Congr. Int. Comm. for Optics: Optics for the Quality of Life, Pts 1 & 2*, G. C. Righini and A. Consortini, Eds., 2003, pp. 771–772.
- [78] H. Minamide and H. Ito, "Coherent terahertz-wave generation and detection over a wide frequency range using DAST crystals," in *Proc. SPIE Nonlinear Freq. Generation and Conversion: Materials, Devices, Appl. VIII*, P. E. Powers, Ed., 2009.
- [79] H. Minamide, J. Zhang, R. Guo, K. Miyamoto, S. Ohno, and H. Ito, "High-sensitivity detection of terahertz wave using nonlinear up-conversion in an organic 4-dimethylamino-N-methyl-4-stilbazolium tosylate crystal," *Appl. Phys. Lett.*, vol. 97, p. 121106(1–3), Sep. 2010.
- [80] M. Tang, H. Minamide, Y. Wang, T. Notake, S. Ohno, and H. Ito, "Tunable terahertz-wave generation from DAST crystal pumped by a monolithic dual-wavelength fiber laser," *Opt. Express*, vol. 19, no. 2, pp. 779–786, Jan. 2011.
- [81] T. Notake, M. Tang, Y. Wang, K. Nawata, H. Ito, and H. Minamide, "Hybrid terahertz-wave source with ultrawideband tunability utilizing organic DAST and BNA crystals," in *2011 Conf. Lasers and Electro-Opt. (CLEO)*, 2011.
- [82] A. Schneider, M. Neis, M. Stillhart, B. Ruiz, R. Khan, and P. Gunter, "Generation of terahertz pulses through optical rectification in organic DAST crystals: Theory and experiment," *J. OSA B—Opt. Phys.*, vol. 23, no. 9, pp. 1822–1835, Sep. 2006.
- [83] K. Reimann, "Table-top sources of ultrashort THz pulses," *Rep. Progr. Phys.*, vol. 70, no. 10, pp. 1597–1632, Oct. 2007.
- [84] K. Nakamura, K. Kasahara, M. Sato, and H. Ito, "Interferometric studies on a diode-pumped Nd:YVO₄ laser with frequency-shifted feedback," *Opt. Commun.*, vol. 121, pp. 137–40, 1995.
- [85] K. Nakamura, F. Abe, K. Kasahara, T. Hara, M. Sato, and H. Ito, "Spectral characteristics of an all solid-state frequency-shifted feedback laser," *IEEE J. Quantum Electron.*, vol. 33, no. 1, pp. 103–111, Jan. 1997.

- [86] K. Kasahara, K. Nakamura, M. Sato, and H. Ito, "Dynamic properties of an all solid-state frequency-shifted feedback laser," *IEEE J. Quantum Electron.*, vol. 34, no. 1, pp. 190–203, Jan. 1998.
- [87] K. Nakamura, T. Miyahara, and H. Ito, "Observation of a highly phase-correlated chirped frequency comb output from a frequency-shifted feedback laser," *Appl. Phys. Lett.*, vol. 72, no. 21, pp. 2631–2633, May 1998.
- [88] K. Nakamura, T. Miyahara, M. Yoshida, T. Hara, and H. Ito, "A new technique of optical ranging by a frequency-shifted feedback laser," *IEEE Photon. Technol. Lett.*, vol. 10, no. 12, pp. 1772–1774, Dec. 1998.
- [89] M. Yoshida, K. Nakamura, and H. Ito, "A new method for measurement of group velocity dispersion of optical fibers by using a frequency-shifted feedback fiber laser," *IEEE Photon. Technol. Lett.*, vol. 13, no. 3, pp. 227–229, Mar. 2001.
- [90] F. V. Kowalski, C. Ndiaye, K. Nakamura, and H. Ito, "Noise waveforms generated by frequency shifted feedback lasers: Application to multiple access communications," *Opt. Commun.*, vol. 231, pp. 149–164, 2004.
- [91] C. Ndiaye, T. Hara, and H. Ito, "Long range, high accuracy optical ranging using frequency-shifted feedback lasers," in *Pacific Rim Conf. Lasers and Electro-Opt.*, Seoul, South Korea, Aug. 26–31, 2007, pp. 419–420.
- [92] Y. Sasaki, Y. Suzuki, K. Suizu, H. Ito, S. Yamaguchi, and M. Imaeda, "Surface-emitted terahertz-wave difference-frequency generation in periodically poled lithium niobate ridge-type waveguide," *Jpn. J. Appl. Phys., Part 2—Lett., Express Lett.*, vol. 45, no. 12–16, pp. L367–L369, Apr. 2006.
- [93] K. Takano, K. Nakagawa, and H. Ito, "Optical tunable delay lines using fiber ring with acousto-optic frequency shifters and EDFAs: I. Experimental demonstration," *IEICE Electron. Express*, vol. 3, no. 20, pp. 442–446, Oct. 2006.
- [94] N. Uesugi, S. Harada, Y. Sasaki, and H. Ito, "Retraction quasi-phase-matching properties of congruent periodically poled LiNbO₃ in type-I second harmonic generation," *J. Appl. Phys.*, vol. 100, no. 8, Oct. 2006.
- [95] T. Yasui, T. Ohtsuka, T. Suzuki, S. Okajima, K. Nakayama, M. Tomioka, K. Kamimura, T. Namekata, H. Minamide, and H. Ito, "Wide range detector using parabolic cylindrical mirror for THz applications," *Int. J. Infrared Millim. Waves*, vol. 27, no. 2, pp. 199–210, Feb. 2006.
- [96] T. Hidaka, A. Ishikawa, J. I. Kojou, T. Ikari, Y. I. Ishikawa, H. Minamide, A. Kudoh, J. I. Nishizawa, and H. Ito, "Adaptive optics instrumentation in submillimeter/terahertz spectroscopy with a flexible polyvinylidene fluoride cladding hollow waveguide," *Review of Scientific Instruments*, vol. 78, no. 8, Aug. 2007.
- [97] R. Guo, S. Ohno, H. Minamide, T. Ikari, and H. Ito, "Highly sensitive coherent detection of terahertz waves at room temperature using a parametric process," *Appl. Phys. Lett.*, vol. 93, no. 2, Jul. 2008.
- [98] M. V. Drummond, J. D. Reis, R. N. Nogueira, P. P. Monteiro, A. L. Teixeira, S. Shinada, N. Wada, and H. Ito, "Error-free wavelength conversion at 160 Gbit/s in PPLN waveguide at room temperature," *Electron. Lett.*, vol. 45, no. 22, pp. 1135–1136, Oct. 2009.
- [99] S. Hayashi, K. Nawata, T. Taira, A. Shikata, K. Kawase, and H. Minamide, "Ultrabright continuously tunable terahertz-wave generation at room temperature," *Sci. Rep.*, vol. 4, no. 5045, p. 5, Jun. 2014.
- [100] F. Qi, S. Fan, T. Notake, K. Nawata, T. Matsukawa, Y. Takida, and H. Minamide, "10 aJ-level sensing of nanosecond pulse below 10 THz by frequency upconversion detection via DAST crystal: More than a 4 K bolometer," *Opt. Lett.*, vol. 39, no. 5, pp. 1294–97, Mar. 2014.
- [101] M. Tang, H. Minamide, Y. Wang, T. Notake, S. Ohno, and H. Ito, "Dual-wavelength single-crystal double-pass KTP optical parametric oscillator and its application in terahertz wave generation," *Opt. Lett.*, vol. 35, no. 10, pp. 1698–1700, May 2010.
- [102] R. Guo, T. Ikari, J. Zhang, H. Minamide, and H. Ito, "Frequency-agile THz-wave generation and detection system using nonlinear frequency conversion at room temperature," *Opt. Express*, vol. 18, no. 16, pp. 16430–16436, Aug. 2010.
- [103] A. Mazhorova, J. F. Gu, A. Dupuis, M. Peccianti, O. Tsuneyuki, R. Morandotti, H. Minamide, M. Tang, Y. Wang, H. Ito, and M. Skorobogatiy, "Composite THz materials using aligned metallic and semiconductor microwires, experiments and interpretation," *Opt. Express*, vol. 18, no. 24, pp. 24632–24647, Nov. 2010.
- [104] Y. Y. Wang, M. Tang, T. Notake, K. Nawata, H. Ito, and H. Minamide, "Sensitive water concentration mapping in thin fresh tissues using tunable THz-wave parametric oscillator," *THz Phys., Devices, Syst. V: Adv. Appl. Industry and Defense*, vol. 8023, 2011.
- [105] M. Takahashi, Y. Ishikawa, and H. Ito, "The dispersion correction and weak-hydrogen-bond network in low-frequency vibration of solid-state salicylic acid," *Chem. Phys. Lett.*, vol. 531, pp. 98–104, Apr. 2012.
- [106] T. Otsuji *et al.*, "Ultrahigh sensitive plasmonic terahertz detector based on an asymmetric dual-grating gate HEMT structure," *Solid-State Electron.*, vol. 78, pp. 109–114, Dec. 2012.
- [107] T. Watanabe, S. B. Tombet, Y. Tanimoto, Y. Wang, H. Minamide, H. Ito, D. Fateev, V. Popov, D. Coquillat, W. Knap, Y. Meziani, and T. Otsuji, "Ultrahigh sensitive plasmonic terahertz detector based on an asymmetric dual-grating gate HEMT structure," *Solid-State Electron.*, vol. 78, pp. 109–114, Dec. 2012.
- [108] S. Boubanga-Tombet, Y. Tanimoto, A. Satou, T. Suemitsu, Y. Wang, H. Minamide, H. Ito, D. V. Fateev, V. V. Popov, and T. Otsuji, "Current-driven detection of terahertz radiation using a dual-grating-gate plasmonic detector," *Appl. Phys. Lett.*, vol. 104, 2014.
- [109] H. Ito, "Breakthroughs in photonics 2013: Terahertz wave photonics," *IEEE Photon. J.*, vol. 6, no. 2, p. 5, Apr. 2014, Art. ID 0701405.

Hiromasa Ito received the bachelor's degree in electrical communication engineering, and the Masters and Ph.D. degrees in electronic engineering all from Tohoku University, Sendai, Japan, in 1966, 1969, and 1972, respectively.

He then joined the Research Institute of Electrical Communication (RIEC) of Tohoku University as a Research Associate. From 1975 to 1976, he was sponsored by the Japan Society for the Promotion of Science (JSPS) as a visiting researcher at Stanford University, Palo Alto, CA, USA, working under noted physicist, Robert Byer. In 1982, he was appointed Associate Professor of Tohoku University in the division of Applied Quantum Electronics at RIEC. He has since been engaged in research on optical coherent waves based on lasers and nonlinear optics, and their applications. In 1993, he became a full professor, and in 1998, he was appointed to the first of two terms (12 years) as Team Leader of the "Tera-Photonics" project of RIKEN, Sendai, part of Japan's major Institute of Physical and Chemical Research network. From 2004 to 2007 he also served as the Director of RIEC. From 2007 to 2010, he was a professor of the Endowed Chair of the University and in 2010 he turned the Tera-Photonics program over to Hiroaki Minamide, becoming a senior scientist at RIKEN. His research interests include lasers, nonlinear optics and nonlinear materials, laser measurements and coherent terahertz science and technology. He pioneered the development and use of the periodically poled lithium niobate and DAST crystals for use as sources and detectors, the coaxial transverse junction surface emitting laser and LED, and most recently the frequency-shifted feedback laser (FSFL), which has already been shown to have a large number of commercial and scientific applications. On the professional engineering front, he was the director of NICHe (New Industry Creation Hatchery Center) of Tohoku University from 2001 to 2003, and the director of Innovation Plaza Miyagi of the JST (Japan Science and Technology Agency) from 2009 to 2012 to work for the promotion of university–industry collaborative research. Starting in 2012 he has been requested to serve as a program officer for a 10-year JST program on "Collaborative Research based on Industrial Demand" for THz technology. He has had a long-term collaboration with Robert Byer at Stanford

University and has been a Visiting Professor at the University of Paris 7 and Dalian Polytechnic University (2007), as well as at Tianjin University in 2010. He has been at Tohoku for more than 50 years and he enjoys running as well as research. It is especially enjoyable in the autumn around RIKEN when he tries to encourage his colleagues to join him.

Prof. Ito is a Fellow of the Optical Society of America (OSA), the Institute of Electronics, Information, and Communication Engineers of Japan (IEICE) and of the Japan Society of Applied Physics (JSAP). He was awarded the IEICE Distinguished Achievement and Contributions Award, and the JSAP Opto/Quantum Electronics Distinguished Achievement Award, for his pioneering work in nonlinear optics, especially his periodically poled LiNbO₃ research. He also received the Kenjiro Sakurai Award of the Optoelectronics and Industry Technology Development Association (OITDA) for his exceptional work in Terahertz wave science and technology.